

ON A MIN-MAX CONJECTURE  
FOR REDUCIBLE DIGRAPHS

Jayme Luiz Szwarcfiter  
Universidade Federal do Rio de Janeiro  
Núcleo Comp. Eletr. e Inst. Matemática  
Caixa Postal 2324  
20.001 - Rio de Janeiro, RJ  
BRASIL

ABSTRACT

A. Frank and A. Gyárfás (1976) have conjectured that in a reducible digraph  $D$  the maximum number of edge disjoint cycles equals the minimum number of edges intersecting all cycles of  $D$ . We prove this conjecture in the special case when  $D$  has at most two distinct dominators. The proof leads to a polynomial time algorithm for finding both the maximum set of cycles and minimum set of edges, in the considered case.

RESUMO

A. Frank e A. Gyárfás (1976) conjecturaram que em um dígrafo redutível  $D$  o número máximo de ciclos disjuntos em arestas é igual ao número mínimo de arestas que interceptam todos os ciclos de  $D$ . Provamos essa conjectura no caso especial em que  $D$  possui no máximo dois denominadores distintos. A prova conduz a um algoritmo polinomial para encontrar tanto o conjunto máximo de ciclos quanto o conjunto mínimo de arestas, no caso considerado.

## 1. Introduction

A conjecture by A. Frank and A. Gyárfás [2] states that the maximum number of edge disjoint cycles of a reducible digraph  $D$  equals the minimum number of edges whose removal turns  $D$  acyclic. In the present paper we prove a special case of this conjecture. The proof is constructive and leads to a polynomial time algorithm for finding such a maximum set of cycles and minimum set of edges, in the considered case.

A flow digraph is a digraph  $D$  together with a vertex  $s \in V(D)$ , called root, such that every vertex of  $D$  is reachable from  $s$ . In particular, if every path from  $s$  to  $v \in V(D)$  contains  $w \in V(D)$  then  $w$  dominates  $v$ . A (fully) reducible digraph is a flow digraph  $D$  such that each cycle  $C$  of  $D$  contains some vertex  $w$  which dominates all the vertices of  $C$ . We call  $w$  a dominator of both  $C$  and  $D$ . See [5,7,9].

Let  $D$  be a general digraph. Denote by

$\alpha_V$  = set of vertex disjoint cycles of  $D$

$\alpha_E$  = set of edge disjoint cycles of  $D$

$\beta_V$  = set of vertices intersecting all cycles of  $D$

$\beta_E$  = set of edges intersecting all cycles of  $D$

Clearly,  $\max |\alpha_V| \leq \min |\beta_V|$  and  $\max |\alpha_E| \leq \min |\beta_E|$ .

$\beta_V$  and  $\beta_E$  are also known as feedback vertex and edge sets, respectively. Recall that the problems of finding the minimum cardinality sets  $\alpha_V$  and  $\alpha_E$  are both NP-hard [3,6].

Theorem 1 (Frank and Gyárfás [2]): If  $D$  is reducible then  $\max |\alpha_V(D)| = \min |\beta_V(D)|$ .

It follows that a minimum feedback vertex set of a reducible digraph can be found in polynomial time [2,4,8].

Conjecture [2]: If  $D$  is reducible then  $\max |\alpha_E(D)| = \min |\beta_E(D)|$ .

We prove this conjecture in the case when  $D$  has at most two distinct dominators.

## 2. The Proof

Throughout this section,  $D$  will always denote a reducible digraph.

Let  $C \equiv v_1, \dots, v_k, v_1$ ,  $k > 1$ , be a cycle of  $D$  and  $v_1$  the dominator of  $C$ . Then edge  $(v_k, v_1)$  is called a back edge of  $D$ .

Lemma 1: Each cycle of  $D$  contains exactly one back edge.

Proof: Let  $v_1, \dots, v_k, v_1$  be a cycle of  $D$  and  $v_1$  its dominator. Then  $(v_k, v_1)$  is a back edge. Suppose that  $C$  contains another back edge  $(v_i, v_{i+1})$ ,  $i \neq k$ . In this case,  $v_{i+1}$  is the dominator of another cycle  $C'$  which contains  $v_i$ . Let  $P$  be a path from  $s$  to  $v_1$  followed by  $v_1, \dots, v_i$ . Then  $P$  meets  $C'$  in a vertex which is not its dominator, a contradiction. That is,  $C$  has exactly one back edge.

Let  $\{r_1, \dots, r_m\}$  be the set of dominators of  $D$ . Denote by  $D^*$  the network obtained by the following construction:

1. Remove all back edges of  $D$ . Let  $D_A$  be the resulting acyclic digraph.

2. Assign a distinct positive label  $x(r_i)$  to each dominator  $r_i$ , such that if  $r_i$  reaches  $r_j$  in  $D_A$  then  $x(r_i) < x(r_j)$ ,  $1 \leq i, j \leq m$  and  $i \neq j$ .

3. For each back edge  $(v, r_j)$  of  $D$ , include a new vertex  $w$  and an edge  $(v, w)$ . Assign to  $w$  the negative label  $-x(r_j)$ .

4. Include the ordered sets

$$S = \{s_1, \dots, s_m\} \quad \text{and} \quad T = \{t_1, \dots, t_m\}$$

of new vertices.  $S$  and  $T$  are the sources and sinks of  $D^*$ , respectively. For  $1 \leq j \leq m$ , include the edge  $(s_j, r_j)$  and an edge to  $t_j$  from every vertex with label  $-x(r_j)$ . Assign to  $s_j$  and  $t_j$  the labels  $x(r_j)$  and  $-x(r_j)$ , respectively.

5. Assign capacity 1 to each edge of  $D_A$  and infinite to those leaving  $S$  and entering  $T$ .

The sets  $S = \{s_i\}$  and  $T = \{t_i\}$  are in normal order if  $x(s_i) < x(s_{i+1})$  and  $x(t_i) > x(t_{i+1})$ ,  $1 \leq i < m$ , respectively.

Lemma 2: Let  $S = \{s_1, \dots, s_m\}$  and  $T = \{t_1, \dots, t_m\}$  be the sources and sinks of  $D^*$ , respectively in normal order. Then there is a one-to-one correspondence between cycles of  $D$  and  $s_j$ - $t_j$  paths of  $D^*$ .

Proof: Let  $v_1, \dots, v_k, v_1$  be a cycle  $C$  of  $D$  and  $v_1$  its dominator. Then  $(v_k, v_1)$  is a back edge and there exists  $w \in V(D^*) - V(D)$  such that  $(v_k, w) \in E(D^*)$  and  $x(w) = -x(v_1)$ . Consequently, for some  $j$ ,  $1 \leq j \leq m$ ,  $(s_j, v_1), (w, t_j) \in E(D^*)$  and therefore  $s_j, v_1, \dots, v_k, w, t_j$  is a path in  $D^*$ . The converse is similar.



Lemma 3: Let  $\{r_1, r_2\}$ ,  $S = \{s_1, s_2\}$  and  $T = \{t_1, t_2\}$  be the dominators, sources and sinks of  $D^*$  in normal order, respectively. Denote by  $f$  a  $S$ - $T$  flow in  $D^*$  having value  $n$  and such that  $f(s_1) \geq f(t_1)$ . Then  $D$  has at least  $n$  edge disjoint cycles,

Proof: Since  $f$  has value  $n$ ,  $D^*$  contains a set  $P$ ,  $|P|=n$ , of  $S$ - $T$  edge disjoint paths. Divide  $P$  into four subsets  $P_1, P_2, P_3$  and  $P_4$ , consisting respectively of  $s_1-t_1$ ,  $s_2-t_2$ ,  $s_1-t_2$  and  $s_2-t_1$  paths. Clearly,  $|P_3| = f(s_1) - |P_1|$  and  $|P_4| = f(t_1) - |P_1|$ . Hence  $|P_3| \geq |P_4|$ . We obtain the required cycles as follows. Each path of  $P_1$  or  $P_2$  corresponds to a cycle of  $D$ , according to lemma 2. Since  $D$  is reducible, each  $s_1-t_2$  path contains  $r_2$ . Consequently, the union of a pair of <sup>paths</sup> cycles, one of  $P_3$  and the other of  $P_4$ , contains two disjoint paths, of types  $s_1-t_1$  and  $s_2-t_2$ , respectively, that is, two new cycles of  $D$ . In addition, the  $|P_3| - |P_4|$  remaining  $s_1-t_2$  paths can be transformed into an equal number of  $s_2-t_2$  paths, by disregarding the  $s_1-r_2$  subpaths and adding edges  $(s_2, r_2)$ . A total of  $n$  edge disjoint cycles of  $D$  has been obtained.

If  $k > 0$  is an integer denote by  $kD^*$  the network obtained from  $D$  as follows:

1. Construct  $D^*$ . Let  $D_1^*, \dots, D_k^*$  be  $k$  identical copies of  $D^*$ , with  $S_i = \{s_{i1}, \dots, s_{im}\}$  and  $T_i = \{t_{i1}, \dots, t_{im}\}$ , respectively the sets of sources and sinks of  $D_i^*$ , in normal order. The vertices of  $S_1$  and  $T_k$  are the sources and sinks of  $kD^*$ , respectively.

2. For  $1 < i < k$  and  $1 < j < m$  include an edge  $(t_{ij}, s_{i+1, j})$  and assign to it infinite capacity.



D\*




Figure 1 : A reducible digraph D and its associated networks.

Lemma 4: If  $kD^*$  has a cut  $X$  of capacity  $n < \infty$  then  $D$  has a feedback edge set  $\beta_E$ , such that  $|\beta_E| \leq n$ .

Proof: Let  $Q = \{e \in E(D) \mid e \in X\}$ . Then  $|Q| \leq n$ . Next, we show that  $Q$  is a feedback edge set. Suppose it is not. Then by lemma 2,  $D^*$  has a path  $p$  of the type  $s_j - t_j$  containing no edge of  $Q$ . By repeating  $p$  in each copy  $D_i^*$  of  $kD^*$  we obtain a  $S_1 - T_k$  path in  $kD^*$  with no edge of  $X$ , a contradiction  $\Delta$

Theorem 2: If  $m \leq 2$ ,  $\max |\alpha_E(D)| = \min |\beta_E(D)|$ .

Proof: If  $m=1$  the theorem follows from lemma 2 and the max-flow min-cut theorem [1] applied to  $D^*$ . When  $m=2$ , construct  $kD^*$ ,  $k = |E(D)|$ . Let  $S_1 = \{s_{11}, s_{12}\}$  and  $T_k = \{t_{k1}, t_{k2}\}$  be the sets of sources and sinks of  $kD^*$ , respectively in normal order, and  $f$  a maximum  $S_1 - T_k$  flow in  $kD^*$ , having value  $n$ . Suppose, initially,  $f(s_{i1}) < f(t_{i1})$ , for all  $1 \leq i \leq k$ . Since  $f(s_{i1}) + f(s_{i2}) = f(t_{i1}) + f(t_{i2}) = n$ , it follows that  $f(s_{i2}) > f(t_{i2})$ ,  $1 \leq i \leq k$ . Because  $f(t_{i2}) = f(s_{i+1,2})$ ,  $1 \leq i < k$ , we conclude that  $f(s_{12}) > f(s_{22}) > \dots > f(s_{k2})$ . However, the latter inequality can not occur because  $f(s_{j2}) < |E(D)|$ ,  $k = |E(D)|$  and all flow values  $f(s_{i2})$  are non negative integers. Consequently, there exists some  $j$ ,  $1 \leq j \leq k$ , such that  $f(s_{j1}) \geq f(t_{j1})$ . Applying lemma 3 to  $D_j^*$ , we conclude that  $D$  contains at least  $n$  edge disjoint cycles, that is,  $\max |\alpha_E(D)| \geq n$ . By the max-flow min-cut theorem,  $kD^*$  has a cut of capacity  $n$  and using lemma 4 it follows that  $D$  contains a feedback edge set of size  $\leq n$ , that is,  $\min |\beta_E(D)| \leq n$ . Hence  $\max |\alpha_E(D)| \geq \min |\beta_E(D)|$ . Since  $\max |\alpha_E(D)| \leq \min |\beta_E(D)|$  the equality follows  $\Delta$

### 3. The Algorithms

The algorithms follow from the proof. Given the reducible digraph  $D$ , construct the network  $kD^*$ ,  $k=|E(D)|$ . Then find the minimum cut  $X$  of  $kD^*$ . The edges of  $D$  which form  $X$  constitute the minimum cardinality feedback edge set  $\beta_E$ . For finding a maximum set of disjoint cycles  $\alpha_E$ , let  $f$  be the maximum S-T flow in  $|E(D)|D^*$ . Next, identify the copy  $D_j^*$  in  $|E(D)|D^*$  such that  $f(s_{j1}) \geq f(t_{j1})$ . Then use the construction of lemma 4 which transforms  $f$  into the desired  $\alpha_E$ .

Both algorithms, for finding a minimum feedback edge set and a maximum set of edge disjoint cycles, have the same complexity as finding a maximum S-T flow in the network  $|E(D)|D^*$ , that is, polynomial in  $|V(D)|$ .

REFERENCES

- [1] L.R.Ford and D.R. Fulkerson, Flows in Networks, Princeton University Press, Princeton, 1962.
- [2] A.Frank and A.Gyárfás, Colloques Internationaux C.N.R.S. Problèmes Combinatoires et Théorie des Graphes (260), 157-158, 1976.
- [3] M.Garey and D.S.Johnson, Computers and Intractability: A Guide to the Theory of NP-Completeness, W.H.Freeman, San Francisco, 1979.
- [4] A.Gyárfás, Blocking the Loops of a Flow-Graph, Zeits. Angew. Math. Mech. 56 (1976), T330-331.
- [5] M.S.Hecht and J.D.Ullman, Characterizations of Reducible Flow Graphs, J. of the ACM 21 (1974), 367-375.
- [6] R.M.Karp, Reducibility among Combinatorial Problems, in R.E. Miller and J.W. Thatcher (eds.), Complexity of Computer Computations, Plenum Press, New York, 85-103, 1972.
- [7] V.N.Kasyanov, Some Properties of Fully Reducible Graphs, Inf. Proc. Letters 2 (1973), 113-117.
- [8] A.Shamir, A Linear Time Algorithm for Finding Minimum Cutsets in Reducible Graphs, SIAM J. Computing 8 (1979), 645-655.
- [9] R.E.Tarjan, Testing Flow Graph Reducibility, J. Comp. Systems Sci. 9 (1974), 355-365.